Supplemental Material

Electron Detachment Dynamics of the Iodide-Guanine Cluster: Does Ionization Occur from the Iodide or from Guanine?

Rosaria Cercola,¹ Kelechi O. Uleanya,¹ and Caroline E. H. Dessent¹*

¹ Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK.

S1 Lowest Energy I⁻⋅G clusters

S2 Collision Induced Dissociation

S1 Lowest Energy I-G clusters

The structures of $I \cdot G$ clusters were studied via DFT, at the B3LYP/6-311++G(2d,2p) level of theory on C, N, O, and H, and 6-311G(d,p) on I with the iodine core electrons being described using the Stuttgart/Dresden (SDD) electron core pseudopotential. 3H, 7H and 9H guanine isomers (with H on G3, G7 or G9) were investigated in their keto and enol tautomeric forms. The enolic forms were studied in their cis and trans isomeric forms (with respect to the -NH₂ group), as described in ref. S1. For each of these structures, different iodine positions near the nucleobase were calculated, and the lowest energy ones are shown in Table S1.

Table S1Calculated structures of the I·G cluster. Calculations were performed were
optimised at the B3LYP/6-311++G(2d,2p) level of theory on C, N, O, and H,
and 6-311G(d,p)/SDD on I.

Structure	Relative Energy ^a (kJ/mol)	Structure	Relative Energy ^a (kJ/mol)	Structure	Relative Energy ^a (kJ/mol)
	0		25.0		33.0
I ⁻ ·G7		I ⁻ ·G9_E_trans		I ⁻ ·G7_E_cis	
	2.43		28.7		38.0
I⁻·G9		I ⁻ ·G9_E_cis		I ⁻ ·G3_E_cis	
	3.45		31.7		40.7
I ⁻ ·G7_E_trans		I ⁻ ·G3_E_trans		I⁻∙G3	

^aRelative energies are zero point corrected energies.

S2 Collision Induced Dissociation

The fragments associated with ground-state thermal fragmentation were identified *via* collision-induced dissociation (CID) in the ion trap of the amaZon SL (Bruker, Daltoniks) [S2]. Here, selected molecular ions are trapped and accelerate to cause collisions with helium buffer gas and thus fragmentation. The acceleration is obtained by applying a voltage to the end-cap electrode, varying its amplitude between 0 and 10% (2.5 V being the maximum). CID was performed on the isolated Γ ·G cluster and the fragment production curves are shown

in Figure S1.




Figure S1. Fragment production curves for I⁻ G upon CID between 0 and 10 % energy.

References

[S1] G. Chunga, H. Oh and D. Lee, J. Mol. Struct. THEOCHEM **730**, 241 (2005).
[S2] J.N. Louris, R.G. Cooks, J.E.P. Syka, P.E. Kelley, G.C. Stafford Jr. and J.F.J. Todd, Anal. Chem. **59**, 1677 (1987).