

Supplemental Table 1. Characteristics of online discussion group participants (n=21 completed surveys/46 participants)

Characteristic	Mean; n(%)
Mean age in years (range)	31 (21-62)
Mean age at diagnosis in years (range)	25 (17-39).
Not diagnosed with endometriosis	1 (5%)
Born in Australia	17 (81%)
Speak a language other than English at home	3 (14%)
Highest level of education	
Secondary school	2 (10%)
Post-secondary trade, certificate, diploma	7 (33%)
Undergraduate degree	5 (24%)
Postgraduate degree	7 (33%)
Worst endometriosis pain experienced (mild-moderate-severe); n=20	18 (90%) severe
Never given birth	17 (81%)
Never tried for > 1 year to conceive without success	17 (81%)

Supplemental Table 2 Characteristics of interview participants

HEALTH PROFESSIONAL #	COUNTRY OF BIRTH	DISCIPLINE	SETTING
1	Australia	General Practitioner (GP)	university health service; ages 17-65; 30% international students all ages
2	Australia	GP	large group practice, mono-cultural
3	Iraq	GP registrar	large regional practice; high patient diversity; especially socioeconomic status (SES)
4	Australia	GP	mid-sized metropolitan clinic; ethnically diverse
5	Malaysia	GP	regional practice, mixed SES
6	UK	GP	group practice; high proportion lower SES
7	Bangladesh	GP	rural setting; multilingual community ; mixed SES
8	Dubai; Indian heritage	GP	private practice; regional city; mixed SES

9	Australia	GP special interest in women's health	metropolitan practice; high SES
10	UK	GP; former paediatrician	regional city; diverse SES and education; culturally diverse including Australian Aboriginal
11	Sweden	GP	regional city; mixed cultural background and SES
12	Australia	Gynaecology; advanced laparoscopy	private specialist practice; inner Melbourne metropolitan
13	England	GP; aboriginal remote community and refugee health experience	Aboriginal Health Service regional city; social welfare support; public chronic pain service; diverse SES