

Paleomagnetism of the Carboniferous Gresford Block, Tamworth Belt, southern New England Orogen: minor counter-clockwise rotation of a primary arc segment

C. Klootwijk*

Research School of Earth Sciences, The Australian National University, Acton, ACT 2601, Australia

*chris.klootwijk@anu.edu.au

SUPPLEMENTARY PAPERS

Australian Journal of Earth Sciences (2020) 67, doi:10.1080/08120099.2020.1674381

Copies of Supplementary Papers may be obtained from the Geological Society of Australia's website (www.gsa.org.au), the Australian Journal of Earth Sciences website (www.ajes.com.au) or from the National Library of Australia's Pandora archive (<https://pandora.nla.gov.au/tep/150555>).

Supplementary paper

Tables (excel workbook)

Table S1. Site and sample details for Gresford Block.

Table S2. Primary and other selected characteristic (*in italics*) high temperature (HT) components (P, P1) in stratigraphic coordinates.

Table S3. Fold test (McFadden, 1990), primary and overprint components (P, P1).

Table S4. Primary and overprint poles (P, P1).

Table S5. Reversal tests for primary and overprint (P, P2) components.

Table S6. Reversal tests for pseudo-primary and overprint (P, P1) components.

Table S7. Low temperature overprint components (L, L1, L2, L3) in geographic coordinates, see Figure 18.

Table S8. High temperature, pre-folding, overprint components (O, P, Q, W) in stratigraphic coordinates.

Table S9. Susceptibility anisotropy in stratigraphic coordinates.

Table S10. Susceptibility anisotropy in geographic coordinates.

Table S11. SNEO pole path data (Figure 17a).