Supplementary material

Figure S1. The micrographs of wound closure of the *in vitro* wound-healing HCEnC model with time after exposure to 0.5, 1, 2, 4 and 8 J/cm² UVA for 2 h. NUVA: without exposure to UVA; 0 to 20h PUVA: at 0-20h post-exposure to UVA for 2 h. Scale bar=50µm.

Figure S2. The micrographs of wound closure with time after exposure to 1 or 2 J/cm² UVA following pretreatment with or without 1.5 mM Asc-2p for 2 h. NUVA: without exposure to UVA; 0 to 20h PUVA: at 0-20h post-exposure to UVA for 2 h. Control: without pretreatment with Asc-2p and exposure to UVA; control+1.5 mM Asc-2p: treatment with 1.5 mM Asc-2p for 2 h without exposure to UVA; 1 J/cm²: exposure to 1 J/cm² for 2 h; 1.5 mM Asc-2p + 1 J/cm²: pretreatment with 1.5 mM Asc-2p for 2 h then exposure to 1 J/cm² UVA for 2 h; 2 J/cm²: exposure to 2 J/cm² UVA for 2 h; 1.5 mM Asc-2p for 2 h then exposure to 2 J/cm² UVA for 2 h; 1.5 mM Asc-2p for 2 h then exposure to 2 J/cm² UVA for 2 h. Scale bar=50μm.