

**Supplemental table 1 Characteristics of participants in 2010 and 2017.**

	2010 cohort	2017 cohort	<i>P</i> value
n	3712	3712	
BMI (kg/m <sup>2</sup> )	23.60±3.61	23.94±3.21	<0.001
SBP (mmHg)	121.07±17.05	125.31±18.27	<0.001
DBP (mmHg)	77.31±11.88	77.70±12.29	<0.001
TC (mmol/L)	4.62±0.88	4.67±0.87	<0.001
TG (mmol/L)	1.05 (0.72-1.61)	1.25 (0.86-1.85)	<0.001
HDL-C (mmol/L)	1.34±0.32	1.30±0.31	<0.001
LDL-C (mmol/L)	2.77±0.75	2.79±0.74	0.164
AIP	-0.07±0.32	0.012±0.32	<0.001
FPG (mmol/L)	5.10±0.52	5.30±0.84	<0.001
UA (mg/dL)	310.88±84.32	340.59±92.58	<0.001
SCr (μmol/L)	65.62±12.68	72.70±14.08	<0.001
eGFR (mL/min/1.73 m <sup>2</sup> )	110.20±10.79	98.11±12.88	<0.001
Hypertension, n (%)	785 (21.1)	1086 (29.3)	<0.001

Data are mean ± SD, median (IQR) or percentage.

BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; TC, total cholesterol; TG, triglyceride; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; AIP, atherogenic index of plasma; FBG, fasting plasma glucose; UA, uric acid; SCr, serum creatinine; eGFR, estimated glomerular filtration rate.