Supplementary Table 1 Pathological features of different histopathological types of anti-GBM disease in 38 patients undergoing renal biopsy

	Focal	Mixed	Crescentic	Sclerotic	p
	(n=6)	(n=5)	(n=22)	(n=5)	
Total glomeruli	18.5 (15.0, 30.5)	13.0 (8.5,	19.0 (15.0, 26.0)	12.0 (10.5, 21.0)	0.355
		24.0)			
Normal glomeruli (%)	71.4 (63.2, 89.6)	17.8 (3.8,	5.8 (0.0, 11.7) ^a	0.0 (0.0, 15.3) ^a	< 0.001
		33.9)			~ 0.001
Global sclerotic	8.5 (0, 13.0)	20.0 (0, 35.0)	0.0 (0, 6.9)	66.7 (55.8,	< 0.001
glomeruli (%)				84.3) ^{a,c}	
Cellular crescents (%)	9.2 ± 10.9	27.4 ± 16.6	$76.7\pm15.6^{a,b}$	11.2 ± 12.7^{c}	< 0.001
Acute tubulointerstitial	1.5 (1.0, 2.3)	3.0 (2.0, 3.0)	3.0 (3.0, 3.0) ^a	3.0 (2.5, 3.0) ^a	< 0.001
lesions score					
Chronic					
tubulointerstitial	1.0 (1.0, 1.0)	2.0 (1.0, 2.0)	1.0 (1.0, 1.0)	2.0 (1.5, 2.0)	0.038
lesions score					
Tubular atrophy and	25.0 (13.8, 32.5)	30.0 (22.5,	22.5 (20.0, 25.0)	50.0 (40.0,	0.002
Interstitial fibrosis (%)		35.0)		50.0) ^{a,c}	

 $^{^{\}mathrm{a}}$ Focal versus mixed, crescentic, and sclerotic (p < 0.05).

 $^{^{}b}$ Mixed versus crescentic and sclerotic (p < 0.05).

 $^{^{\}circ}$ Crescentic versus sclerotic (p < 0.05).