

Supplementary Table A. Mean (M) and standard deviation (SD) trials for each condition included in the analyses in TS patients and control group.

Condition	TS Patients M \pm SD	Control group M \pm SD
Anger cue	40 \pm 8	39 \pm 8
Happiness cue	35 \pm 9	39 \pm 10
Sadness cue	37 \pm 12	38 \pm 9
Neutral cue	36 \pm 11	38 \pm 12
Congruent target linked to Anger cue	32 \pm 7	33 \pm 6
Congruent target linked to Happiness cue	28 \pm 7	31 \pm 8
Congruent target linked to Sadness cue	29 \pm 10	31 \pm 8
Congruent target linked to Neutral cue	30 \pm 9	31 \pm 8
Incongruent target linked to Anger cue	7 \pm 3	6 \pm 2
Incongruent target linked to Happiness cue	7 \pm 2	8 \pm 3
Incongruent target linked to Sadness cue	8 \pm 3	7 \pm 2
Incongruent target linked to Neutral cue	6 \pm 2	7 \pm 3
No-response target linked to Anger cue	10 \pm 3	11 \pm 2
No-response target linked to Happiness cue	9 \pm 3	11 \pm 2
No-response target linked to Sadness cue	10 \pm 3	10 \pm 2
No-response target linked to Neutral cue	9 \pm 3	10 \pm 3