

Table 1 Supplementary. Baseline characteristics and risk factors for preterm birth in women with singleton pregnancies undergoing cerclage, based on Body Mass Index (BMI).

Variables	Overall	BMI Classification			
		Normal (n = 16)	Overweight (n = 35)	Obese (n = 44)	<i>p</i>
Maternal age (years)	30.1 ± 0.7	30.1 ± 2.1	29.7 ± 1.1	30.5 ± 0.9	0.85
Race/Ethnicity					<0.01
Black	48 (50.5)	4 (25.0)	12 (34.3)	32 (72.7)	
Hispanic	25 (26.3)	4 (25.0)	12 (34.3)	9 (20.5)	
White	12 (12.6)	6 (37.5)	4 (11.4)	2 (4.6)	
Other	10 (10.5)	2 (12.5)	7 (20.0)	1 (2.3)	
Marital Status					0.48
Married	45 (47.4)	8 (50.0)	19 (54.3)	18 (40.9)	
Other	50 (52.6)	8 (50.0)	16 (45.7)	26 (59.1)	
Primigravida	11 (11.6)	1 (6.3)	5 (14.3)	5 (11.4)	0.71
Previous preterm birth	60 (63.2)	13 (81.3)	22 (62.9)	25 (56.8)	0.22
Previous loss at 16 to 24 weeks	58 (63.7)	8 (50.0)	21 (60.0)	29 (72.5) (n=40)	0.24
Previous cone biopsy	13 (13.8)	2 (12.5)	8 (23.5) (n=34)	3 (6.8)	0.1
Previous cerclage	26 (27.7)	6 (37.5)	10 (28.6)	10 (23.3) (n=43)	0.55
Gestational age at cerclage (weeks)	16.5 ± 0.3	15.9 ± 0.6	16.6 ± 0.6	16.7 ± 0.5	0.66
Cervical length < 25 mm prior to cerclage	30 (40.0)	2 (14.3) (n=14)	10 (41.7) (n=24)	18 (48.7) (n=37)	0.08

Data are presented as mean ± SE and number (percent) where applicable.