

Table S2. Results from the analysis of variance of linear mixed-effects models. We tested the effect of the factors Snowmelt (levels: Early, Mid, Late), Site (levels: North, South) and Year (levels: 2014, 2015) and their interactions on the timing of reproductive phenophases. The plot from which an observation was made was the random effect. The left-hand column lists the species and phenophases which were analysed and the factor levels which were included in each model. Each model was fitted twice with one of two response variables: either the number of days after snowmelt, or the day of year, when a phenophase occurred for the first time. Non-significant ($p>0.05$) interactions or main effects were identified by the maximum likelihood ratio test and excluded from the model. For each main effect and each significant interaction, degrees of freedom (in subscript), F -values and p -values are shown.

Phenophase and species	Days after snowmelt				Day of year			
	Snowmelt	Site	Year	Significant interactions	Snowmelt	Site	Year	Significant interactions
Flower opening <i>Empetrum nigrum</i> North, Early-Mid, 2014-2015	$F_{1,6}=38.0668$, $p=0.0008$	-	$F_{1,103}=9.0706$, $p=0.0033$	Snowmelt x Year ($F_{1,103}=408.6525$, $p<.0001$)	$F_{1,6}=16.8$, $p=0.0064$	-	$F_{1,103}=58.314$, $p<.0001$	Snowmelt x Year ($F_{1,103}=231.645$, $p<.0001$)
North-South, Early, 2015	-	$F_{1,5}=3.5734$, n.s.	-	-	-	$F_{1,5}=29.79$, $p=0.0028$	-	-
<i>Phyllodoce caerulea</i> North-South, Late, 2014-2015	-	n.s.	$F_{1,183}=1251.818$, $p<.0001$	n.s.	-	$F_{1,6}=77.9$, $p=0.0001$	$F_{1,182}=441.8$, $p<.0001$	Site x Year ($F_{1,182}=145.6$, $p<.0001$)
South, Mid-Late, 2014-2015	$F_{1,4}=67.7635$, $p=0.0012$	-	$F_{1,136}=596.6245$, $p<.0001$	Snowmelt x Year ($F_{1,136}=4.4209$, $p=0.0373$)	$F_{1,4}=76.92$, $p=0.0009$	-	$F_{1,136}=18.74$, $p<.0001$	Snowmelt x Year ($F_{1,136}=9.83$, $p=0.0021$)
<i>Vaccinium myrtillus</i>	Could not be tested				Could not be tested			
<i>Vaccinium uliginosum</i> North-South, Early, 2014- 2015	-	$F_{1,5}=12.8045$, $p=0.0159$	$F_{1,104}=140.5085$, $p<.0001$	Site x Year ($F_{1,104}=80.4184$, $p<.0001$)	-	$F_{1,5}=0.33$, n.s.	$F_{1,104}=211.505$, $p<.0001$	Site x Year ($F_{1,104}=8.134$, $p=0.0052$)
South, Early-Mid, 2014-2015	$F_{1,5}=27.6181$, $p=0.0033$	-	$F_{1,103}=387.4231$, $p<.0001$	Snowmelt x Year ($F_{1,103}=21.556$, $p<.0001$)	$F_{1,5}=5.204$, n.s.	-	$F_{1,103}=166.945$, $p<.0001$	Snowmelt x Year ($F_{1,103}=25.483$, $p<.0001$)

Phenophase and species	Days after snowmelt				Day of year			
	Snowmelt	Site	Year	Significant interactions	Snowmelt	Site	Year	Significant interactions
South, Early-Mid-Late, 2015	$F_{2,6}=13.4142$, $p=0.0061$	-	-	-	$F_{2,6}=17.61$, $p=0.0031$	-	-	-
<i>Vaccinium vitis-idaea</i> North-South, Early-Mid, 2014-2015	$F_{1,10}=194.217$, $p<.0001$	$F_{1,10}=37.311$, $p=0.0001$	$F_{1,122}=1122.112$, $p<.0001$	Snowmelt x Site ($F_{1,10}=46.273$, $p<.0001$) Site x Year ($F_{1,122}=16.754$, $p=0.0001$) Snowmelt x Site x Year ($F_{1,122}=9.407$, $p=0.0027$)	$F_{1,10}=85.3$, $p<.0001$	$F_{1,10}=41.3$, $p=0.0001$	$F_{1,125}=776.4$, $p<.0001$	Snowmelt x Site ($F_{1,10}=65.6$, $p<.0001$)
North-South, Early-Mid-Late, 2015	$F_{2,13}=22.1912$, $p=0.0001$	$F_{1,13}=17.4129$, $p=0.0011$	-	n.s.	$F_{2,14}=11.409$, $p=0.0011$	n.s	-	n.s
Flower senescence <i>Empetrum nigrum</i> North, Early-Mid, 2014-2015	$F_{1,6}=6.89187$, $p=0.0393$	-	$F_{1,249}=0.3435$, n.s.	Snowmelt x Year ($F_{1,249}=156.0788$, $p<.0001$)	$F_{1,6}=18.991$, $p=0.0048$	-	$F_{1,249}=52.111$, $p<.0001$	Snowmelt x Year ($F_{1,249}=97.623$, $p<.0001$)
North-South, Early, 2014- 2015	-	$F_{1,6}=1.943$, n.s.	$F_{1,194}=0.3946$, n.s.	Site x Year ($F_{1,194}=175.0225$, $p<.0001$)	-	$F_{1,6}=38.86$, $p=0.0008$	$F_{1,194}=72.73$, $p<.0001$	Site x Year ($F_{1,194}=81.12$, $p<.0001$)
North-South, Early-Mid, 2015	$F_{1,10}=3.4952$, n.s.	$F_{1,103}=166.945$, n.s.	-	Snowmelt x Site ($F_{1,10}=13.6157$, $p=0.0042$)	$F_{1,10}=90.96$, $p<.0001$	$F_{1,10}=59.74$, $p<.0001$	-	Snowmelt x Site ($F_{1,10}=10.05$, $p=0.01$)
<i>Phyllodoce caerulea</i> North-South, Late, 2014-2015	-	$F_{1,6}=6.8521$, $p=0.0397$	$F_{1,199}=1233.784$, $p<.0001$	n.s.	-	$F_{1,6}=0$, n.s.	$F_{1,198}=416.47$, $p<.0001$	Site x Year ($F_{1,198}=178.32$, $p<.0001$)
South, Mid-Late, 2014-2015	$F_{1,4}=92.2543$, $p=0.0007$	-	$F_{1,141}=600.4432$, $p<.0001$	Snowmelt x Year ($F_{1,141}=4.7627$, $p=0.0307$)	$F_{1,4}=9.73$, $p=0.0356$	-	$F_{1,141}=63.907$, $p<.0001$	Snowmelt x Year ($F_{1,141}=8.484$, $p=0.0042$)
<i>Vaccinium myrtillus</i>	Could not be tested				Could not be tested			

Phenophase and species	Days after snowmelt				Day of year			
	Snowmelt	Site	Year	Significant interactions	Snowmelt	Site	Year	Significant interactions
<i>Vaccinium uliginosum</i>								
South, Early-Mid-Late, 2014-2015	$F_{1,6}=23.3715$, $p=0.0015$	-	$F_{1,80}=323.4421$, $p<.0001$	Snowmelt x Year ($F_{1,73}=10.418$, $p=0.0001$)	$F_{2,6}=16.61$, $p=0.0036$	-	$F_{1,80}=127.17$, $p<.0001$	Snowmelt x Year ($F_{2,80}=6.34$, $p=0.0028$)
North-South, Early, 2014-2015	-	$F_{1,5}=3.1984$, n.s.	$F_{1,85}=137.9683$, $p<.0001$	Site x Year ($F_{1,85}=53.4447$, $p<.0001$)	-	n.s.	$F_{1,86}=200.736$, $p<.0001$	n.s.
<i>Vaccinium vitis-idaea</i>								
North-South, Early-Mid, 2014-2015	$F_{1,9}=27.954$, $p=0.0005$	$F_{1,9}=0.2471$, n.s.	$F_{1,80}=160.0848$, $p<.0001$	Snowmelt x Year ($F_{1,80}=4.5655$, $p=0.0357$) Site x Year ($F_{1,80}=10.6739$, $p=0.0016$) Snowmelt x Site x Year ($F_{1,80}=7.3165$, $p=0.0083$)	n.s.	n.s.	$F_{1,83}=134.761$, $p<.0001$	n.s.
North-South, Early-Mid-Late, 2015	$F_{2,9}=5.9101$, $p=0.023$	$F_{1,9}=5.2323$, $p=0.048$	n.s.	n.s.	n.s.	n.s.	-	n.s.
Fruitset visible								
<i>Empetrum nigrum</i>								
North, Early-Mid, 2014-2015	$F_{1,6}=38.153$, $p=0.0008$	-	$F_{1,136}=8.665$, $p=0.0038$	n.s.	$F_{1,6}=26.979$, $p=0.002$	-	$F_{1,143}=7.71$, $p=0.0062$	n.s.
North-South, Early, 2014-2015	-	$F_{1,6}=1.0266$, n.s.	$F_{1,132}=47.6274$, $p<.0001$	Site x Year ($F_{1,132}=74.0540$, $p<.0001$)	-	$F_{1,6}=36.93$, $p=0.0009$	n.s.	n.s.
North-South, Early-Mid, 2015	$F_{1,180}=4.0784$, n.s.	$F_{1,9}=0.6976$, n.s.	-	Snowmelt x Site ($F_{1,9}=7.3917$, $p=0.0237$)	$F_{1,9}=62.04$, $p<.0001$	$F_{1,9}=65.24$, $p<.0001$	n.s.	Snowmelt x Site ($F_{1,9}=12.45$, $p=0.0064$)
<i>Phyllodoce caerulea</i>								
North-South, Late, 2014-2015	-	n.s.	$F_{1,168}=227.94$, $p<.0001$	n.s.	-	$F_{1,6}=4.06$, n.s.	$F_{1,167}=2.82$, n.s.	Site x Year ($F_{1,167}=60.8$, $p<.0001$)
South, Mid-Late, 2014-2015	$F_{1,4}=108.6893$, $p=0.0005$	-	$F_{1,97}=95.8318$, $p<.0001$	Snowmelt x Year ($F_{1,97}=6.7793$, $p=0.0107$)	Could not be tested			
<i>Vaccinium myrtillus</i>								
North-South, Late, 2014	-	n.s.	-	-	-	n.s.	-	-

Phenophase and species	Days after snowmelt				Day of year			
	Snowmelt	Site	Year	Significant interactions	Snowmelt	Site	Year	Significant interactions
North, Late, 2014-2015	-	-	$F_{1,62}=200.809$, $p<.0001$	-	-	-	$F_{1,59}=103.35$, $p<.0001$	-
<i>Vaccinium uliginosum</i> South, Early-Mid-Late, 2015	$F_{2,6}=13.3311$, $p=0.0062$	-	-	-	$F_{2,6}=10.05$, $p=0.0121$	-	-	-
North-South, Early, 2014-2015	Could not be tested				-	$F_{1,5}=12.6$, $p=0.0164$	$F_{1,86}=182.254$, $p<.0001$	n.s.
South, Early-Mid, 2014-2015	Could not be tested				n.s.	-	$F_{1,89}=94.5$, $p<.0001$	n.s.
<i>Vaccinium vitis-idaea</i> North, Mid, 2014-2015	-	-	$F_{1,28}=153.426$, $p<.0001$		-	-	$F_{1,28}=104.11$, $p<.0001$	-
North-South, Early-Mid-Late, 2015	$F_{2,11}=15.2962$, $p=0.0007$	$F_{1,11}=20.1995$, $p=0.0009$	-	n.s.	$F_{2,9}=4.909$, $p=0.0362$	$F_{1,9}=0.198$, n.s.	-	Snowmelt x Site ($F_{2,9}=2.879$, n.s.)
South, Early-Mid, 2014-2015	$F_{1,4}=3.2173$, n.s.	-	$F_{1,43}=447.7688$, $p<.0001$	Snowmelt x Year ($F_{1,43}=8.8636$, $p=0.0048$)	$F_{1,4}=22.976$, $p=0.0087$	-	$F_{1,44}=124.603$, $p<.0001$	n.s.
Fruit ripe <i>Empetrum nigrum</i> North, Early-Mid, 2014-2015	$F_{1,4}=24.2883$, $p=0.0079$	-	$F_{1,33}=185.0477$, $p<.0001$	Snowmelt x Year ($F_{1,33}=18.3001$, $p=0.0002$)	n.s.	-	$F_{1,34}=332.32$, $p<.0001$	n.s.
North-South, Early, 2014-2015	-	$F_{1,4}=1.544$, n.s.	$F_{1,74}=605.9165$, $p<.0001$	Site x Year ($F_{1,74}=137.9141$, $p<.0001$)	-	$F_{1,4}=16.42$, $p=0.0154$	$F_{1,75}=702.07$, $p<.0001$	n.s.
North-South, Early-Mid, 2015	$F_{1,7}=14.063$, $p=0.0072$	$F_{1,7}=9.9051$, $p=0.0162$	-	n.s.	n.s.	$F_{1,8}=6.3$, $p=0.0363$	-	n.s.
<i>Vaccinium myrtillus</i> North-South, Late, 2014	-	n.s.	-	-	-	n.s.	-	-

Phenophase and species	Days after snowmelt				Day of year			
	Snowmelt	Site	Year	Significant interactions	Snowmelt	Site	Year	Significant interactions
<i>Vaccinium uliginosum</i> South, Early-Mid, 2014	Could not be tested				Could not be tested			

"-" – This term was not part of the analysis

"Could not be tested" – The data did not fulfil the ANOVA assumptions