


**Supplemental Figure 1.** Spectrogram showing the broadband pulse candidate call type (2,500-point fast Fourier transform, Hann window with 50% overlap). Calls were recorded in Cormorant Channel, British Columbia, Canada, during a single occasion when minke whales were the only baleen whale visually observed. Spectrogram on the right shows a higher resolution view of the frequency range contained in the red box in the spectrogram on the left.



**Supplemental Figure 2.** Spectrogram showing the tonal waver candidate call type (2,500-point fast Fourier transform, Hann window with 50% overlap). Calls were recorded in Cormorant Channel, British Columbia, Canada, during a single occasion when minke whales were the only baleen whale visually observed.

**Supplemental Table 1.** Summary of acoustic parameters describing two call types recorded in the presence of minke whales in Cormorant Channel, British Columbia. All calls of each type were heard on a single occasion, and no other baleen whales were observed in the study area at these times. All statistics are reported as mean  $\pm$  standard deviation.

<b>Call Type</b>	<b>Sample Size</b>	<b>Low Frequency (Hz)</b>	<b>High Frequency (Hz)</b>	<b>Duration (ms)</b>	<b>Peak Frequency (Hz)</b>
Broadband Pulse	2	40 $\pm$ 8	2007 $\pm$ 62	190 $\pm$ 71	234 $\pm$ 44
Tonal Waver	22	106 $\pm$ 21	156 $\pm$ 7	940 $\pm$ 309	131 $\pm$ 14