

Activity of hydrothermal fluid at the bottom of a lake and its influence on the development of high-quality source rocks: Triassic Yanchang Formation, southern Ordos Basin, China

J. You^{a,b}, Y. Liu^a, D. Zhou^c, Q. Zheng^b, K. Vasichenko^d and Z. Chen^e

^aState Key Laboratory of Continental Dynamics, Department of Geology, Northwest University, Xi'an 710069, China; ^bSchool of Energy Engineering, Yulin University, Yulin 719000, China; ^cCollege of Earth Science and Engineering, Shandong University of Science and Technology, Qingdao 266510, China; ^dCollege of Economics and Management, Northwest A&F University, Yangling 712100, China; ^eSchool of Earth Sciences and Engineering, Xi'an Petroleum University, Xi'an 710065, China

First author: You Jiyuan Email: 2373375337@qq.com

Corresponding author: Liu Yiqun liu-yiqun@263.com

SUPPLEMENTARY PAPERS

Australian Journal of Earth Sciences (2019) 66, [doi:10.1080/08120099.2019.1612783](https://doi.org/10.1080/08120099.2019.1612783)

Copies of Supplementary Papers may be obtained from the Geological Society of Australia's website (www.gsa.org.au), the Australian Journal of Earth Sciences website (www.ajes.com.au) or from the National Library of Australia's Pandora archive (<https://pandora.nla.gov.au/tep/150555>).

Supplementary paper

Table S1: Major element and rare earth element concentrations from the Bawangzhuang profile, Z233 well, and H269 well. Fe₂O₃ = total iron. (excel file)