Supplemental file 1

Calculation of low field effect (LFE) by two-proton model

Here the Hamiltonian below is assumed $(I_1 = I_2 = 1/2)$.

$$
\hat{H} = \sum_{i=1,2} \frac{\mathcal{S}_i \mu_B B_0}{\hbar} \hat{S}_{Z_i} + a_1 \hat{S}_1 \bullet \hat{I}_1 + a_2 \hat{S}_2 \bullet \hat{I}_2 \approx \omega_0 \left(\hat{S}_{Z_1} + \hat{S}_{Z_2} \right) + a_1 \hat{S}_1 \bullet \hat{I}_1 + a_2 \hat{S}_2 \bullet \hat{I}_2 \tag{1}
$$

Difference in g values are ignored because we will discuss the spin dynamics at very low magnetic fields. Out of $2^{2+2} = 16$ eigenstates of the Hamiltonian, the 7 states below are involved in the LFE as Lewis et al.[1] reported (for the full description, see the supporting information for the original paper).

$$
|2\rangle = c_{2+} |T_{+1}\alpha\beta\rangle + \frac{c_{2-}}{\sqrt{2}} (|T_0\alpha\alpha\rangle + |S\alpha\alpha\rangle),
$$

\n
$$
|4\rangle = c_{1+} |T_{+1}\beta\alpha\rangle - \frac{c_{1+}}{\sqrt{2}} (|T_0\alpha\alpha\rangle - |S\alpha\alpha\rangle),
$$

\n
$$
|6\rangle = \frac{1}{\sqrt{2}} (|T_0\alpha\beta\rangle + |S\alpha\beta\rangle),
$$

\n
$$
|7\rangle = c_{1+}c_{2+} |T_{+1}\beta\beta\rangle + \frac{c_{1+}c_{2-}}{\sqrt{2}} (|T_0\beta\alpha\rangle + |S\beta\alpha\rangle)
$$

\n
$$
+ \frac{c_{1-}c_{2+}}{\sqrt{2}} (|T_0\alpha\beta\rangle - |S\alpha\beta\rangle) + c_{1-}c_{2-} |T_{-1}\alpha\alpha\rangle,
$$

\n
$$
|11\rangle = \frac{1}{\sqrt{2}} (|T_0\beta\alpha\rangle - |S\beta\alpha\rangle),
$$

\n
$$
|12\rangle = \frac{c_{2+}}{\sqrt{2}} (|T_0\beta\beta\rangle - |S\beta\beta\rangle) + c_{2-} |T_{-1}\beta\alpha\rangle,
$$

\n
$$
|14\rangle = \frac{c_{1+}}{\sqrt{2}} (|T_0\beta\beta\rangle + |S\beta\beta\rangle) + c_{1-} |T_{-1}\alpha\beta\rangle,
$$

\n
$$
|F = -1
$$

\n
$$
|14\rangle = \frac{c_{1+}}{\sqrt{2}} (|T_0\beta\beta\rangle + |S\beta\beta\rangle) + c_{1-} |T_{-1}\alpha\beta\rangle,
$$

\n(2)

and

$$
\omega_{2} = \frac{\omega_{0} + \mu_{2}}{2} + \frac{a_{1} - a_{2}}{4},
$$
\n
$$
\omega_{4} = \frac{\omega_{0} + \mu_{1}}{2} + \frac{-a_{1} + a_{2}}{4},
$$
\n
$$
\omega_{6} = \frac{a_{1} + a_{2}}{4},
$$
\n
$$
\omega_{7} = \frac{\mu_{1} + \mu_{2}}{2} - \frac{a_{1} + a_{2}}{4},
$$
\n
$$
\omega_{11} = \frac{a_{1} + a_{2}}{4},
$$
\n
$$
\omega_{12} = -\frac{\omega_{0} - \mu_{2}}{2} + \frac{a_{1} - a_{2}}{4},
$$
\n
$$
\omega_{13} = -\frac{\omega_{0} + \mu_{1}}{2} - \frac{a_{1} - a_{2}}{4},
$$
\n
$$
\omega_{14} = -\frac{\omega_{0} + \mu_{1}}{2} - \frac{a_{1} - a_{2}}{4},
$$
\n
$$
\omega_{15} = -1
$$

where

$$
\mu_i = \sqrt{\omega_0^2 + a_i^2} \tag{4}
$$

and

$$
c_{i\pm}^2 = \frac{\mu_i \pm \omega_0}{2\mu_i} \tag{5}
$$

The time dependent singlet population for the singlet-born radical pair is described as

$$
P_{\rm s}(t) = \frac{1}{Z} \sum_{m,n} \left| \left\langle n \right| \hat{P}_{\rm s} \left| m \right\rangle \right|^2 e^{i\omega_{mn}t} \tag{6}
$$

By applying a magnetic field, breakdown of zero-field degeneracy occurs for the 4 pairs of eigenstates (note that $\omega_6 = \omega_{11}$ at any magnetic field), so that the oscillation becomes active as

$$
P_{\rm S}^{2-4}(t) = |\langle 2|\hat{P}_{\rm S}|4\rangle|^2 e^{i\omega_{2-4}t} = \frac{1}{4}c_{1-}^2c_{2-}^2 \exp\left\{i\left(-\frac{\mu_1 - \mu_2}{2} + \frac{a_1 - a_2}{2}\right)t\right\}
$$

\n
$$
P_{\rm S}^{6-7}(t) = |\langle 6|\hat{P}_{\rm S}|7\rangle|^2 e^{i\omega_{6-7}t} = \frac{1}{4}c_{1-}^2c_{2+}^2 \exp\left\{-i\left(\frac{\mu_1 + \mu_2}{2} - \frac{a_1 + a_2}{2}\right)t\right\}
$$

\n
$$
P_{\rm S}^{7-11}(t) = |\langle 7|\hat{P}_{\rm S}|11\rangle|^2 e^{i\omega_{7-1}t} = \frac{1}{4}c_{1+}^2c_{2-}^2 \exp\left\{i\left(\frac{\mu_1 + \mu_2}{2} - \frac{a_1 + a_2}{2}\right)t\right\}
$$

\n
$$
P_{\rm S}^{12-14}(t) = |\langle 12|\hat{P}_{\rm S}|14\rangle|^2 e^{i\omega_{7-1}t} = \frac{1}{4}c_{1+}^2c_{2+}^2 \exp\left\{i\left(-\frac{\mu_1 - \mu_2}{2} + \frac{a_1 - a_2}{2}\right)t\right\}
$$
 (7)

Let us assume that the total HF coupling is constant $(a_1 > a_2 > 0)$ as

$$
a \equiv a_1 + a_2
$$

\n
$$
r \equiv \frac{a_2}{a_1} \quad (0 \le r \le 1)
$$

\n
$$
R \equiv \frac{a_0}{a} \quad (0 \le R)
$$

\n(8)

Thus

$$
a_1 = \frac{a}{1+r}
$$
\n
$$
a_2 = \frac{ar}{1+r}
$$
\n
$$
\mu_1 = \sqrt{a_0^2 + a_1^2} = \sqrt{a^2 R^2 + \frac{a^2}{(1+r)^2}} = a \sqrt{R^2 + \frac{1}{(1+r)^2}}
$$
\n
$$
\mu_2 = \sqrt{a_0^2 + a_2^2} = \sqrt{a^2 R^2 + \frac{a^2 r^2}{(1+r)^2}} = a \sqrt{R^2 + \frac{r^2}{(1+r)^2}}
$$
\n(10)

The angular frequency parts of eq. 7 is thus rewritten as

$$
\omega_{2-4} = \omega_{12-14} = \left(-\frac{\mu_1 - \mu_2}{2} + \frac{a_1 - a_2}{2}\right) = \frac{a}{2} \left(-\sqrt{R^2 + \frac{1}{(1+r)^2}} + \sqrt{R^2 + \frac{r^2}{(1+r)^2}} + \frac{1-r}{1+r}\right) \ge 0
$$
\n
$$
-\omega_{6-7} = \omega_{7-11} = \left(\frac{\mu_1 + \mu_2}{2} - \frac{a_1 + a_2}{2}\right) = \frac{a}{2} \left(\sqrt{R^2 + \frac{1}{(1+r)^2}} + \sqrt{R^2 + \frac{r^2}{(1+r)^2}} - 1\right) \ge 0
$$
\n(11)

These indicate that the mixing frequencies linearly increases with the total coupling *a* at very low magnetic fields where ω < α ($R \sim 0$). The first derivative of ω ₋₄ is

$$
\frac{\partial \omega_{2-4}}{\partial r} = \frac{a}{2} \left(\frac{1}{\sqrt{\left(1+r\right)^4 + R^2 \left(1+r\right)^6}} + \frac{r}{\sqrt{r^2 \left(1+r\right)^4 + R^2 \left(1+r\right)^6}} - \frac{2}{\left(1+r\right)^2} \right) \tag{12}
$$

This value is maximised when $R = 0$ irrespective of r values, so that

$$
\frac{\partial \omega_{2-4}}{\partial r}(\text{max}) = \frac{a}{2} \left(\frac{1}{\sqrt{(1+r)^4}} + \frac{r}{\sqrt{r^2 (1+r)^4}} - \frac{2}{(1+r)^2} \right)
$$

=
$$
\frac{a}{2} \left(\frac{1}{(1+r)^2} + \frac{1}{(1+r)^2} - \frac{2}{(1+r)^2} \right) = 0
$$
 (13)

Thus

$$
\frac{\partial \omega_{2-4}}{\partial r} \le 0 \tag{14}
$$

at any *r* and *R* values in the defined range.

On the other hand, the first derivative of $-\omega_{6-7}$ is

$$
\frac{-\partial \omega_{6-7}}{\partial r} = \frac{a}{2} \left(\frac{r}{\sqrt{r^2 (1+r)^4 + R^2 (1+r)^6}} - \frac{1}{\sqrt{(1+r)^4 + R^2 (1+r)^6}} \right)
$$
\n
$$
= \frac{a}{2} \left(\frac{r}{\sqrt{r^2 (1+r)^4 + R^2 (1+r)^6}} - \frac{r}{\sqrt{r^2 (1+r)^4 + r^2 R^2 (1+r)^6}} \right)
$$
\n(15)

Since $0 < r < 1$, $R^2 > r^2 R^2$, so that

$$
\frac{-\partial \omega_{6-7}}{\partial r} \le 0\tag{16}
$$

From these calculations, it turns out that all the four angular frequencies monotonically decrease with increasing *r*, which means that the mixing frequency is lower for similar HF coupling constants for the two radicals.

Dependence of ω_{2-4} and $-\omega_{6-7}$ on *r* at $R = 0.168$, which is the same condition for the $B_0 = 0.6$ mT and $a/2\pi = 100$ MHz as shown in Figure 3 in the main text, is shown in Figure S1-1.

Figure S1-1. Dependence of the relative mixing frequency $\omega(a_1 + a_2)$ on the relative HF coupling a_2/a_1 at a low magnetic field of $a_0/a = 0.168$.

The mixing frequencies are almost ten times lower than the total HF coupling at $r = 0$ (HF coupling on only one radical) and reduces significantly with increasing *r* (more balanced HF couplings for the two radicals).

[1] A. M. Lewis, T. P. Fay, D. E. Manolopoulos, C. Kerpal, S. Richert and C. R. Timmel, J. Chem. Phys. **149**, 034103 (2018).