Supplemental Table 2. Changes (post- minus pre-treatment) in sIL-1R1 and sIL-1R2 levels of individual patients were correlated with changes (post- minus pre-treatment) in levels of different targets and values for visual parameters with correlation coefficients r>0.5. Targets with p<0.05 are highlighted in bold.

Changes in sIL-1R1 Correlated with		
Changes in Target Levels	Pearson Correlation Coefficient (p-value)	Spearman Correlation Coefficient (p-value)
sIL-1R2	0.733 (0.015)	0.551 (0.098)
sRAGE	0.698 (0.024)	0.539 (0.107)
IL-17A	0.669 (0.034)	0.563 (0.089)
IL-18	0.567 (0.087)	0.515 (0.127)
VEGF-D	-0.572 (0.083)	-0.539 (0.107)
Changes in Vision Parameters	Pearson Correlation Coefficient (p-value)	Spearman Correlation Coefficient (p-value)
СМТ	0.236 (0.511)	0.054 (0.881)
VA	-0.178 (0.622)	-0.139 (0.700)
Changes in sIL-1R2 Correlated with		
Changes in Target Levels	Pearson Correlation Coefficient (p-value)	Spearman Correlation Coefficient (p-value)
sIL-1R1	0.733 (0.015)	0.551 (0.098)
IL-17A	0.733 (0.015)	0.781 (0.007)
VEGF-D	-0.722 (0.018)	-0.636 (0.047)
Changes in Vision Parameters	Pearson Correlation Coefficient (p-value)	Spearman Correlation Coefficient (p-value)
CMT	0.191 (0.595)	0.224 (0.533)
VA	0.078 (0.829)	0.194 (0.590)