

Dual-target anti-Alzheimer's disease agents with both iron ion chelating and monoamine oxidase-B inhibitory activity

Zhisheng Mi ^{a, #}, Bing Gan ^{a, b, #}, Sihang Yu ^{c, #}, Jianan Guo ^a, Changjun Zhang ^a,
Xiaoying Jiang ^a, Tao Zhou^d, Jing Su ^{d, *}, Renren Bai ^{a, *}, Yuanyuan Xie ^{a, e, *}

^aCollege of Pharmaceutical Science, Zhejiang University of Technology, Hangzhou, China

^bGuiyang Institute for Food and Drug Control, Guiyang, China

^cDepartment of Pathophysiology, College of Basic Medical Sciences, Jilin University, Changchun, China

^dSchool of Food Science and Biotechnology, Zhejiang Gongshang University, Hangzhou, China

^eCollaborative Innovation Center of Yangtze River Delta Region Green Pharmaceuticals, Zhejiang University of Technology, Hangzhou, China

[#]These authors contributed equally to this work



Correspondence:

Dr. Yuanyuan Xie, Key Laboratory for Green Pharmaceutical Technologies and Related Equipment of Ministry of Education, College of Pharmaceutical Sciences, Zhejiang University of Technology, Hangzhou 310014, China. Email: xyycz@zjut.edu.cn



Dr. Renren Bai, College of Pharmaceutical Sciences, Zhejiang University of Technology, Hangzhou 310014, China. Email: renrenbai@zjut.edu.cn

Dr. Jing Su, Department of Pathophysiology, College of Basic Medical Sciences, Jilin University, Changchun, China. Email: sujing@jlu.edu.cn



27a





27b





27c





27d





27e





27f





27g





27h



27i



27j



27k

