

**Halogenated boroxine dipotassium trioxohydroxytetrafluorotriborate $K_2[B_3O_3F_4OH]$
inhibits emerging multidrug-resistant and β -lactamase-producing opportunistic pathogens**

Ana Maravić^{a*}, Tomislav Rončević^b, Lucija Krce^b, Nada Ilić^b, Borivoj Galić^c, Vedrana Čikeš Čulić^d, Ivana Carev^e

^a Department of Biology, Faculty of Science, University of Split, Ruđera Boškovića 33, Split, Croatia.

^b Department of Physics, Faculty of Science, University of Split, Ruđera Boškovića 33, Split, Croatia.

^c (†) Faculty of Natural Sciences, University of Sarajevo, Bosnia and Herzegovina.

^d Department of Medicinal Chemistry and Biochemistry, Medical School, University of Split, Šoltanska 2, 21000 Split, Croatia.

^e Department of Biochemistry, Faculty of Chemistry and Engineering, University of Split, Ruđera Boškovića 35, 21000 Split, Croatia.

*** Corresponding author:** Ana Maravić, Department of Biology, Faculty of Science, University of Split, Ruđera Boškovića 33, 21000 Split, Croatia. Tel.: +38521619281, Fax: +38521619227, E-mail: amaravic@pmfst.hr




Figure S1. The permeability of the membrane of boroxine treated *S. aureus* ATCC 29213. Percentage of PI positive cells measured by flow cytometry after incubation of bacterial cells with 1024 µg/ml (-----) and 16384 µg/ml (——) boroxine is shown.