

Supplemental Table 1: Exogenous Endophthalmitis Survey Questions:

1. Please specify the range of the approximate number of exogenous endophthalmitis patients seen at your institution per year.
 - a. 0-50
 - b. 51-100
 - c. 101-150
 - d. 151-200
 - e. 201+
 - f. Do not know
2. Do you give oral antibiotics for exogenous endophthalmitis?
 - a. Yes
 - b. No
 - c. Do not know
3. Please specify the name (s) and duration of oral antibiotic therapy. (Ex/moxifloxacin for 5 days).
4. Please specify name(s) of intravitreal antibiotics routinely administered for endophthalmitis at your institution. (Ex/vancomycin and ceftazidime).
5. What is the preferred method for lowering IOP during the administration of an intravitreal injection?
 - a. Vitreous tap pre-injection
 - b. Vitreous tap post-injection
 - c. Anterior chamber paracentesis pre-injection
 - d. Anterior chamber paracentesis post-injection
 - e. Other, please specify in box below
 - f. Do not know
6. Are intravitreal steroids routinely administered with intravitreal antibiotic injection in bacterial endophthalmitis?
 - a. Yes
 - b. No
 - c. Do not know
7. Please list the indications for performing pars plana vitrectomy (PPV) for exogenous endophthalmitis at your institution.
8. What is the preferred method for obtaining cultures in exogenous endophthalmitis patients?
 - a. Vitreous tap

- b. Anterior chamber paracentesis
 - c. Vitrectomy
 - d. Other, please specify in box below
- 9. What is the routine ED disposition for exogenous endophthalmitis patients?
 - a. Treated and discharged
 - b. Admitted to hospital
 - c. Other, please specify in box below
 - d. Do not know